

<https://observatoiredeparis.psl.eu/salt-on-io.html>



Salt on Io

Date de mise en ligne : mercredi 1er janvier 2003

**Observatoire de Paris - PSL Centre de recherche en astronomie et
astrophysique**

A team of French and American astronomers, led by E. Lellouch, from Paris-Meudon Observatory, has observed the presence of salt (NaCl) in Jupiter's moon Io's tenuous atmosphere. This species is most probably produced through volcanic emission. Its presence provides an explanation to the "clouds" of atomic sodium that have been observed to surround Io for almost 30 years.

Figure 1. Two views of Jupiter's satellite Io and its environment by the Galileo spacecraft. Left : Color mosaic showing the global aspect of Io's surface and several manifestations of its volcanic activity. Note the presence of two volcanic plumes, one at the limb (Pillan, 140 km altitude), and the other near the center (Prometheus, seen from above). <http://photojournal.jpl.nasa.gov/catalog/PIA01081> Right : A view of the atomic sodium cloud scattering solar light at 589 nm. The very bright spot on Io is the Prometheus plume, also seen is scattered light.

<http://photojournal.jpl.nasa.gov/catalog/PIA00593> The atmosphere of Jupiter's moon Io is one of the most peculiar of the Solar System. In 1979, the Voyager spacecraft revealed active volcanism (Figure 1, left) at the surface of the satellite and discovered a local, tenuous SO₂ atmosphere. Since 1990, millimeter-wave observations acquired at IRAM (French-German-Spanish telescope) and UV observations with HST provided a somewhat more detailed description of this atmosphere. The typical surface pressure is about 1 nanobar, and, in a unique fashion in the Solar System, the atmosphere exhibits strong horizontal variations, being apparently concentrated in an equatorial band. The main atmospheric compounds are SO₂, SO and S₂. The atmosphere is probably produced, on the one hand by direct volcanic output, and on the other hand by the sublimation of SO₂ ices that cover Io's surface. However, it has been long suspected that Io's atmosphere must contain other chemical species. As early as 1974, visible imaging and spectroscopy revealed a "cloud" of atomic sodium (Figure 1, right), roughly centered about Io's orbit. Detailed subsequent studies of this cloud indicated a complex structure, including notably "fast sodium" features, for the production of which the role of molecular ions (NaX⁺) was evidenced. These discoveries naturally raised the question of the origin of sodium in Io's environment. From the brightness of the optical emissions of Na, one can estimate that about 10²⁶-10²⁷ sodium atoms leave Io each second. In 1999, chlorine in atomic and ionized form was discovered around Io, with an abundance comparable to that of sodium (while the cosmochemical abundance of Na is about 15 times that of Cl). This suggests a common origin, NaCl being a natural plausible parent of both. At the same time, on the basis of thermochemical equilibrium calculations, NaCl was proposed to be an important compound of Io's volcanic magmas, with an abundance relative to SO₂ as high as several percent.

Figure 2. The spectral lines of the salt molecule NaCl at frequencies of 143 and 234 GHz detected with the 30m telescope of IRAM on 15-17 January 2002. The spectral resolution is 40 kHz. Based on these discoveries and predictions, an observing campaign was conducted by E. Lellouch, from Paris Observatory, and several French and American colleagues at the IRAM 30-m radiotelescope in January 2002. Two rotational lines of NaCl at 143 and 234 GHz were unambiguously detected (Figure 2.). Because the vapor pressure of this salt is entirely negligible, NaCl cannot be in sublimation equilibrium with Io's surface and its presence must directly result from continuous volcanic output. It appears to be a minor atmospheric species. The most plausible physical model depicts the NaCl atmosphere as more localized than SO₂, due to its very short lifetime (a few hours at most), and probably restricted to the volcanic centers. The local NaCl abundance in this model is 0.3-1.3 % of SO₂, significantly lower than predicted. From the line strengths, volcanic emission rates of (2-8)x10²⁸ NaCl molecules per second can be derived. According to photochemical and escape models, only a small fraction of these molecules escape from Io (about 0.1 %). A somewhat larger amount (1-2 %) leaves Io in atomic form after being photolyzed to Na and Cl. The vast majority of the volcanically-emitted NaCl molecules fall back to the surface where they condense out, potentially contributing to the white color of some of Io's terrains. In conclusion, it appears that NaCl provides an important source of sodium and chlorine in Io's environment ; however the precise chemical nature of the NaX⁺ molecular ions remains to be elucidated.

Reference :

E. Lellouch, G. Paubert, J.I. Moses, N.M. Schneider, and D.F. Strobel. "Volcanically-emitted sodium chloride as a source for Io's neutral clouds and plasma torus" *Nature*, 2 January 2003, 421, p. 45-47

Contact

- Emmanuel Lellouch (Observatoire de Paris, LESIA)